

Some Statistics to Chew On...

- Head and neck squamous cell carcinoma HNSCC make up about 4% of all cancers in the United States.
- It is MUCH more common in China/Southeast Asia
- $^{\circ}$ More than 70% of squamous cell carcinoma of the head and neck are estimated to be avoidable by lifestyle changes
- HPV has been established as a causative agent in the oropharyngeal squamous cell carcinoma and biologically active HPV can act as a prognosticator with better overall survival than HPV-negative tumors.

 $https://oraleancerfoundation.org/wp-content/uploads/2016/03/nasopharynx.pdf \\ https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6031231/ \\ https://www.cancer.gov/types/head-and-neck/head-neck-fact-sheet#q4$


Oral Cavity Cancer Epidemiology of Oral Cancer


- 95% SCCA
- Incidence 6 per 100,000
- Average age 60 years
- Tobacco and ETOH synergistic
- Betel chewing
- Poor oral hygiene




UNC Tobacco Treatment Program • If you smoke or use tobacco and are considering becoming tobacco free, contact the UNC Tobacco Treatment Program for support! We work with everyone: patients, community members and loved ones. • https://www.med.unc.edu/fammed/fammedcenter/tobacco-treatment-program/thinking-of-quitting/









Treatment for Head and Neck Cancers

1.Surgery
2.Radiation
3.Systemic Therapy
(chemotherapy, targeted therapy, immunotherapy)









